


# MATHEMATICS SEMINAR

## PEAKS AND DESCENTS OF PERMUTATIONS

Given a permutation  $\pi = \pi_1\pi_2\cdots\pi_n \in \mathfrak{S}_n$ , we say an index  $i$  is a peak if  $\pi_{i-1} < \pi_i > \pi_{i+1}$ . Let  $P(\pi)$  denote the set of peaks of  $\pi$ . Given any set  $S$  of positive integers, define  $P_S(n) = \{\pi \in \mathfrak{S}_n : P(\pi) = S\}$ . In 2013, Billey, Burdzy, and Sagan showed that for all fixed subsets of positive integers  $S$  and sufficiently large  $n$ ,  $|P_S(n)| = p_S(n)2^{n-|S|-1}$  for some polynomial  $p_S(x)$  depending on  $S$ . They conjectured that the coefficients of  $p_S(x)$  expanded in a binomial coefficient basis centered at  $\max(S)$  are all nonnegative. In this talk, we will share a recursive formula for  $p_S(n)$  we use to prove that their “positivity conjecture” is true. It remains an open question to find a combinatorial meaning of these non-negative coefficients. Near the end of the talk, we will discuss various current developments regarding this topic, including some similar questions replacing “peaks” by “descents.” No prerequisites.

**Friday, March 24<sup>th</sup>**  
**3:30 pm – 4:30 pm**  
**Seidler Hall 220**



**Alexander Diaz-Lopez, Ph.D.**  
Visiting Assistant Professor at Swarthmore College  
2016-2018 Associate Editor for Notices of the AMS

REFRESHMENTS SPONSORED BY

